



Original article

Impact of measuring pneumonia severity index (PSI) in the management of community acquired pneumonia in Emergency Department

Eduardo Esteban-Zubero^{1*}  , Marta Rubio-Gómez² , Alberto García-Noain³ , Beatriz Sierra-Bergua³ ,
Joaquín Gómez-Bitrian³  and Francisco José Ruiz-Ruiz³ 

¹Emergency Department, Hospital San Pedro, Logroño, Spain, ²Emergency Department, Hospital Universitario Miguel Servet, Zaragoza, Spain, ³Emergency Department, Hospital Clínico Universitario Lozano Blesa, Zaragoza, Spain

*Author to whom correspondence should be addressed

Received: 03-11-2022, Revised: 26-11-2022, Accepted: 07-12-2022, Published: 31-12-2022

Copyright © 2022 Esteban-Zubero et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Esteban-Zubero et al. (2022) Impact of measuring pneumonia severity index (PSI) in the management of community acquired pneumonia in Emergency Department. *Mediterr J Pharm Pharm Sci.* 2 (4): 70 - 78.
<https://doi.org/10.5281/zenodo.7479792>.

Keywords: Pneumonia, pneumonia severity index, Emergency Department, Spain

Abstract: Community acquired pneumonia is a prevalent disease in the Emergency Department (ED). The literature reveals that clinical practice could not be related with guidelines recommendations. The aim of this study is to determine the impact of the implementation of the recommendations of the Spanish society of Emergency Medicine in the Hospital Clínico Universitario Lozano Blesa (Zaragoza) ED in the management of community acquired pneumonia. Use of Pneumonia Severity Index (PSI) estimation was used to assess the adherence. This study was carried out from December, 2014 to February, 2015. Data was compared with the previous two months. The indicators management (discharge or admission decision, adequacy and anti-biotherapy) as well as the incidence of PSI estimation before and after the intervention in these parameters were evaluated. 209 patients were included, 97 before the intervention and 112 after the intervention. No significant differences were observed in the calculation of PSI. A significant decrease of admissions was observed after the intervention in the patients in whom the PSI was calculated (68.8% vs. 45.0%, $p < 0.05$). A greater use of the Observation Room was aimed at those patients in whom the PSI was calculated (06.3% vs. 17.5%, $p < 0.05$). PSI calculation significantly increased antibiotic prescription adherence (88.9% vs. 75.2%, $p < 0.05$). There were no modifications in the prescription after the intervention. In conclusion, PSI is a useful and effective measure to achieve a greater adherence to the recommendations. However, despite the positive trend in the use of the PSI and its interpretation, a low-intensity intervention is not sufficient to generalize its use.

Introduction

Community-acquired pneumonia (CAP) is the leading cause of death from infectious disease in developed countries (10.0% to 14.0%, depending on age and associated risk factors) and is the origin of a large part of complications admitted in the

Emergency Department (ED), including sepsis and septic shock [1, 2]. In addition, pneumonia is the third leading reason for hospital admission accounting for 544 000 hospitalizations from the ED annually [3]. These rates are extremely

important due to the great variability existing between different centers and among clinicians in the approach and management of all the diagnostic-therapeutic process of patients with CAP [4 - 6]. This fact may be one of the reasons which explain the existence of different hospital admission rates, the choice of antibiotic regimen, the intensity of care or use of resources [7]. This variability implies that the prognosis and evolution of patients with CAP also be different [8]. It is estimated that 75.0% of all CAP are attended in the ED, which reveals the important role of the emergency physician in the initial management of these processes [1, 3, 7]. The decisions to be made by the emergency physician are critical, including the requirement of hospital admission, the appropriate location and the care that the patients require. All of them determine the prognosis of the patient, the requirement of laboratory tests and microbiological studies, the election of antimicrobial agent, intensity of clinical observation and the use of socio-sanitary resources [7]. All these measurements are related with the final costs, which increase from 08 to 25 times in the patient admitted in the hospital compared with the discharge from the ED [9]. Along with the decision of the patient's destination, the early administration of the antimicrobial agent and hemodynamic and respiratory support measures are the most important factors in the evolution and mortality of patients with CAP [10]. Due to that, in recent years several authors are developing protocols or clinical practice guidelines (CPG) to manage these patients in the ED, especially in critical patients [11 - 14]. Most of them have been performed by specialists in respiratory medicine, analyzing only the patients admitted to the hospital, although others have recently been published that include patients from the ED [15, 16]. While the effectiveness and efficiency of the CPG is recognized, it is observed a low rate of adherence, and up to 35.0% - 65.0% confess do not use them [17, 18]. Due to that, the correct implementation of CPG in the ED agreed with the rest of specialists and adapted to the center is probably the main tool to decrease clinical variability and improve process management [15, 16]. The aim of this study was to determine the adherence in the management of ED

in the CAP healthcare process and the impact of a low intensity intervention attending the recommendations of the Spanish Society of Emergency Medicine [18] about the CAP healthcare process. Pneumonia Severity Index (PSI) estimation was used to assess the adherence. Discharge or admission decision, adequacy and anti-biotherapy will be the indicators management compared before and after the intervention.

Materials and methods

The study was carried out in the Hospital Clínico Universitario Lozano Blesa (Zaragoza). It is a third-level center with 800 beds belonging to the Aragon Health Service (SALUD) with a reference area of 275 000 inhabitants. The ED has an internal medicine area attended by emergency physicians of its own staff and residents of different medical specialties.

Study design: Observational, single-blind study with prospective follow-up of patients in two phases: before and after an intervention consisting of the implementation of the recommendations of the Spanish Society of EM [19] together with training sessions on it to all emergency physicians and residents of the center.

Study period and studied population: The study was carried out from December 1st, 2014 to February 1st, 2015. Before this period (November 2014), the training and different sessions about the recommendations were realized in the ED. To compare the results of the intervention, data was compared with a retrospective cohort of the previous two months (September 25th, 2014 to November 25th, 2014). To be included, the patients had to meet the following criteria: be adults (≥ 18 years old) and diagnosed with CAP in the ED by their responsible doctors. Immunosuppressed patients and patients hospitalized in the previous two weeks were excluded. Subjects in which the responsible doctor did not maintain the final diagnosis of CAP after 30 days (codes 481, 482, 483, 485, 486 and 507 of the International Classification of Diseases, Ninth Revision, Clinical Modification [ICD-9-MC]) and to those who had a final diagnosis of a different etiology were also

excluded. Nor were patients who were diagnosed with a second episode of CAP during the study period included in the study. The study met the ethical standards of the hospital and was approved by the Clinical Research Ethics Committee of Hospital Clinico Universitario Lozano Blesa (Zarazona) Hospital (2014). The follow-up of all the patients was carried out through the computerized clinical history of the ED, Hospital and Primary Care.

Study intervention: Between November 25th and 30th, 2014, the recommendations of the Spanish Society of Emergency Medicine [19] was delivered complete and in a diptych, and training sessions on it were given to all emergency physicians and residents of the center, asking them to systematically apply the recommendations from now on. To evaluate and analyze the performance of the emergency physician in each case, it was defined as "adequate management" when there was agreement with the measures and treatments carried out by them with the aforementioned guide

[19] and "inadequate management" when it did not coincide. This assessment was carried out independently for the request of complementary studies in the ED (laboratory, microbiological and biomarkers), prognostic assessment and decision of the patient's destination according to the PSI and for the choice and administration of treatment in the ED. To assess the prognosis and severity of patients with CAP and the admission decision, the PSI was used, whose risk classes were used according to the original authors' proposals [20], although different additional criteria were included according to the recommendations of the Spanish Society of Emergency Medicine, the main representation of EM in our area. These additional criteria were also explained and transmitted in the training sessions. Thus, it was considered that all the patients belonging to risk classes IV-V (PSI) and those belonging to risk classes I-III should be admitted to the ward in the presence of one or more of the risk factors or additional criteria resumed in **Table 1**.

Table 1: Additional criteria and risk factors that determine the admission of patients

<p>PaO₂ < 60 mmHg or O₂ saturation by pulse oximetry < 90%.</p> <p>Evidence of uncompensated comorbidity.</p> <p>Pleural effusion (encapsulated, ≥ 2 cm on chest radiograph in the lateral decubitus position).</p> <p>Multilobar or bilateral radiological involvement.</p> <p>Criteria for severe sepsis or septic shock.</p> <p>High probability or suspicion of bacteremia due to the clinical situation and / or biomarkers such as PCR > 90 mg/ml and/or procalcitonin > 1 ng / ml.</p> <p>Situations or factors that prevent correct home treatment such as oral intolerance, social problems. (Dependent patient without available caregiver, psychiatric disorders, alcoholism, etc.).</p> <p>Lack of response to previous antibiotic treatment (after 72 h of starting adequate antibiotic treatment in the presence of clinical or radiological deterioration).</p>
--

Variables collected: The patients were divided into the pre-intervention group (Pre-I) and post-intervention group (Post-I). All the clinical, exploratory, radiological, analytical, patient destination (discharge, hospital admission, Observation Room (OBS) in the ED admission and sociodemographic variables included in the PSI [20] were collected. 20 variables establish the index, providing a score according to their summation. As a result, patients may be classified into five categories or classes (I-V) based on 30-day mortality. Thus, classes I-III group patients

with mild CAP and a low risk of mortality (< 03.0%), class IV includes patients with an intermediate risk of dying (08 - 10%), while class V is made up of patients with a high risk of dying (27.0 -31.0%). According to this classification, discharge from the ED and ambulatory treatment is recommended in classes I and II, unless there is hypoxemia. It is recommended admission to short-stay observation units in class III and hospital admission in classes IV and V. The value and grade of PSI was calculated in patients in whom it had not been collected.

Statistical analysis

Mean and standard deviation (SD), range, median and percentage, as appropriate, were used to describe the demographic, clinical, evolutionary and treatment characteristics of the patients in both phases. The comparison of percentage between phases (pre and post implantation of the CPG) was performed using the Chi-square or Fisher's exact test, as appropriate, and that of scale and ordinal variables, using the Student's *t*-test and Mann-Whitney *U*-test as applicable. The statistical analysis was performed with the SPSS-15 program for Windows and value of $p < 0.05$ was considered significant; all the contrasts were bilateral.


Results

In this study, 209 patients were diagnosed with CAP, 97 in the pre-intervention group (pre-I) and 112 in the post-intervention group (post-I). The mean age was 69.5 ± 19.8 years and the distribution by sex showed 56.6% of men ($n = 118$) and 43.5% of women ($n = 91$). PSI was calculated in the pre-I group in the 32.0% of the patients ($n = 31$) whereas in the post-I in 35.7% ($n = 40$), thus being calculated globally in 34.0% ($n = 71$). There were no statistically significant differences between the two groups ($p > 0.05$). Likewise, it is appreciated that in the retrospective calculation, more than half of the patients ($n = 57$, 58.8%) in the pre-I group belonged to assessment groups IV and V, which are indicative of hospital admission. These values are similar to the post-I group ($n = 54$ patients, 48.2%). Adherence to the recommendations attending PSI calculation and patients' destination was assessed before and after the intervention. Comparing the number of admissions, a decrease was observed in

the patients in whom the PSI was calculated, this decrease being greater after the intervention (68.8% pre-I vs. 45.0% post-I, $p < 0.05$) (**Figure 1**). Regarding the use of the OBS, a greater use of the OBS after the intervention was aimed at those patients in whom the PSI was calculated (06.3% pre-I vs. 17.5% post-I, $p < 0.05$) (**Figure 1**). The total breakdown of the patients at the different PSI levels as well as their destination (home/admission and OBS) may be observed in **Table 2**.

Attending to antibiotic treatment, the most prescribed antibiotic was levofloxacin ($n = 120$, 57.4% of the total), followed by amoxicillin-clavulanic ($n = 36$, 17.2%), levofloxacin + ceftriaxone ($n = 9$, 04.3%), moxifloxacin ($n = 09$, 04.3%), ceftriaxone + azithromycin ($n = 08$, 03.8%), imipenem ($n = 05$, 02.4%), amoxicillin-clavulanic + azithromycin ($n = 05$, 2.4%), cefditoren ($n = 01$, 00.5%), and others ($n = 16$, 07.7%). The most frequent lack of adherence was the non-association of macrolide with beta-lactam in admitted patients (12.9%). As previously observed in patients' destination where PSI was calculated, in those patients in whom the PSI was determined, greater adherence to the recommendations was observed (88.9% vs. 75.2%, $p < 0.05$) (**Figure 2**). There were no changes in the prescription after the intervention performed (80.4% of adequacy in pre-I and 79.5% in post-I, $p > 0.05$). All this data may be observed in **Table 2**. A decrease of hospital admissions was observed in the patients in whom the PSI was calculated after the intervention. In addition, a greater use of the OBS after was aimed at those patients in whom the PSI was calculated.

Figure 1: Patient's PSI score attending the area of admission prior and after intervention.




PSI: Pneumonia Severity Index, OBS: Observation Unit, Pre-I: Pre-intervention group, Post-I: Post-intervention group. * $p < 0.05$.

Table 2: Patients distribution and destination attending to their PSI index

<i>Patients destination after ED (Home/Admission) attending to PSI evaluation or not in the ED</i>						
<i>PSI no calculated</i>						
<i>PSI value</i>	I n = 19	II n = 16	III n = 22	IV n = 61	V n = 20	Total n = 138
<i>Home</i>	16	06	04	04	00	29
<i>n (%)</i>	84.2	37.5	18.2	6.6	0.0	21
<i>Admission</i>	03	18	18	57	20	109
<i>n (%)</i>	(15.8%)	(81.8%)	(81.8%)	(93.4%)	(100%)	(71%)
<i>PSI calculated</i>						
<i>PSI value</i>	I n = 16	II n = 10	III n = 15	IV n = 26	V n = 04	Total n = 71
<i>Home</i>	15	09	08	00	00	32
<i>n (%)</i>	(93.7%)	(90.0%)	(53.3%)	(00.0%)	(00.0%)	(45.1%)
<i>Admission</i>	01	01	07	26	04	39
<i>n (%)</i>	(06.3%)	(10.0%)	(46.7%)	(100%)	(100%)	(54.9%)
<i>Patients distribution attending to PSI</i>						
<i>Pre-I</i>	I n = 12 (12.4%)	II n = 12 (12.4%)	III n = 16 (16.5%)	IV n = 45 (46.3%)	V n = 12 (12.4%)	Total n = 97
<i>Post-I</i>	I n = 23 (20.5%)	II n = 14 (12.5%)	III n = 21 (18.8%)	IV n = 42 (37.5%)	V n = 12 (10.7%)	Total n = 112
<i>Total</i>	I n = 35 (16.8%)	II n = 26 (12.4%)	III n = 37 (17.7%)	IV n = 87 (41.6%)	V n = 24 (11.5%)	Total n = 209
<i>Patients destination (Home/Admission) after the attendance in the ED</i>						
<i>Pre-I group</i>						
<i>PSI value</i>	I n = 12	II n = 12	III n = 16	IV n = 45	V n = 12	Total n = 97
<i>Home</i>	n = 11 (91.7%)	n = 05 (41.7%)	n = 05 (31.3%)	n = 04 (08.9%)	n = 00 (00%)	n = 24 (24.7%)
<i>Admission</i>	01 (08.3%)	07 (58.3%)	11 (68.7%)	41 (91.1%)	12 (100%)	73 (75.3%)
<i>Post-I group</i>						
<i>PSI value</i>	I n = 23	II n = 14	III n = 21	IV n = 42	V n = 12	Total n = 112
<i>Home</i>	20 (87.0%)	10 (71.4%)	07 (33.3%)	00 (00.0%)	0 (00.0%)	37 (33.0%)
<i>Admission</i>	03 (13.0%)	04 (28.6%)	14 (66.7%)	42 (100%)	12 (100%)	75 (67.0%)
<i>Total</i>						
<i>PSI value</i>	I (n = 35)	II (n = 26)	III (n = 37)	IV (n = 87)	V (n = 24)	Total (n = 209)
<i>Home</i>	31 (88.6%)	15 (57.7%)	12 (32.4%)	04 (04.6%)	00 (00.0%)	61 (29.2%)
<i>Admission</i>	04 (11.4%)	11 (42.3%)	25 (67.6%)	83 (95.4%)	12 (100%)	148 (70.8%)
<i>Use of Observation Room (OBS)</i>						
<i>PSI no calculated</i>	Pre-I (n = 92, 94.8%)		Post-I (n = 103, 92%)		Total (n = 195, 93.3%)	
<i>PSI calculated</i>	Pre-I (n = 05, 5.2%)		Post-I (n = 09, 8%)		Total (n = 14, 6.7%)	
<i>Adequacy of antibiotic treatment attending to PSI evaluation or not</i>						
<i>PSI no calculated</i>	No adequate (n = 34, 28.8%)		Adequate (n = 103, 75.2%)		Total (n = 137, 100%)	
<i>PSI calculated</i>	No adequate (n = 08, 11.1%)		Adequate (n = 64, 88.9%)*		Total (n = 72, 100%)	
<i>Total</i>	No adequate (n = 42, 20.1%)		Adequate (n = 167, 79.9%)		Total (n = 209, 100%)	
<i>Adequacy of antibiotic treatment</i>						
<i>Pre-I</i>	No adequate (n = 19, 19.6%)		Adequate (n = 78, 80.4%)		Total (n = 97, 100%)	
<i>Post-I</i>	No adequate (n = 23, 20.5%)		Adequate (n = 89, 79.5%)		Total (n = 112, 100%)	
<i>Total</i>	No adequate (n = 42, 20.1%)		Adequate (n = 167, 79.9%)		Total (n = 209, 100%)	

PaO2: Partial pressure of oxygen in arterial blood; O2: Oxygen; PCR: C-reactive protein.

Figure 2: Antibiotic adherence attending the measurement or not of PSI score.



In those patients in whom the PSI was calculated, a greater adherence to the antibiotic recommendations prescription was observed. PSI: Pneumonia Severity Index. * $p < 0.05$.

Discussion

The implementation of a CPG in an ED that includes biomarkers of inflammation and infection as well as comorbidities and physical exploratory signs may become an effective tool to decrease the variability management of CAP. It includes the request for analytical and microbiological studies, the administration of the appropriate early treatment and the final destination of the patient. In addition, it improves the overall mortality of the process, decrease the incidence of improper discharges and inadequate admissions (and its consequences) and reduce the time necessary to achieve clinical stabilization and, consequently, decrease hospital stay. PSI estimation increases the adherence to CPG, especially attending to patient's destination as well as antibiotic treatment, as we observed. The intervention did not evaluate mortality rates as well as hospital stay, being one limitation of the study in the measure of the impact of PSI evaluation. However, after our brief intervention, we observed that the PSI calculation decreased number of hospital admissions, increased admission in OBS and the adherence to recommendations of GPC was higher. It has previously been observed that utilization of CAP management guidelines may improve the clinical outcomes [11, 21, 22]. However, while the benefits of adherence to CAP management guidelines are

clear and accepted universally as a goal in CAP management, strategies aimed at guideline implementation have met with mixed success [23, 24]. Our adherence ratio to the recommendations of CAP management (32.0% in pre-I group, 35.7% in post-I group, 35% globally) is similar to the previous published data [1, 25].

The decision of hospital admission of the patient or not will determine the request for complementary studies and the type of treatment and, consequently, the prognosis and evolution of the patient. Due to that, it is important to reduce improper registrations and unjustified admissions. It is observed in the literature that the routine implantation of a CPG in an ED that incorporates the PSI causes a decrease in the percentage of unjustified admissions, in addition to a greater adequacy of the place of treatment as well as the adequacy and precocity of antibiotic treatment [15, 26] as we observed. Our results revealed a significant decrease of admissions in the patients in whom the PSI was calculated, this decrease being greater after the intervention (68.8% pre-I vs. 45.0% post-I, $p < 0.05$). In addition, it is known that the variability of the percentage of admissions between different centers and doctors is very large when a CPG is not followed, and even when adherence is high, given the existence of criteria or reasons not contemplated in the PSI [4, 5]. Some examples are

observed in the literature. In an observational study carried out in 12 ED with high follow-up of a CPG with the PSI scale, the authors observed that 37.4% of the patients with PSI I-III were admitted and 03.2% of the cases with PSI IV-V were discharged [18]. These authors were in line with the previous studies where the 44.7% of low-risk patients (PSI I-III) were admitted, and of these, 20.0% had no justification or any criteria for doing so. In another recent study [27], to assess the adequacy of hospital admissions for CAP, a lower proportion of inadequate admissions (06.07%) was found in the low-risk group (PSI I-III) but a large proportion of undesirable discharge (according to the PSI and additional criteria), which reached 32.89%. As we commented previously, we observed a decrease in the global admission rate from 68.8% to 45.0% after the intervention in the patients in whom the PSI was calculated. This result was statistically significant, observing a trend of adequacy to recommendations if PSI was calculated, especially in PSI scores of I-III. However, this data was not correlated with the intervention in a general manner, being not observed any statistical differences excepting the increased use of OBS. Our results agree with the previous studies, where the estimation of PSI observed a non-adherence to the GPC [27]. However, our inadequacy was

mostly due to inappropriate admissions other than inappropriate admissions of discharges with PSI I-III. These results could be explained due to the existence of clinical, social and capacity aspects for oral treatment that are not recognized in the PSI and sometimes justify admission [28, 29]. The antibiotic adherence also increased in a significant manner in our study if PSI was calculated. However, we do not observe any differences after the intervention. As previous results observed in our research, it may be due to the low strength of the intervention. However, these results agree with the previous data, being observed an increased adherence to recommendations and validation of PSI to antibiotic prescription, especially if a multi-faceted education intervention is performed [30].

Conclusion: The PSI calculation is confirmed as a useful and effective measure to achieve greater adherence to the CPG. Likewise, the PSI calculation improves adherence to empirical treatment recommendations in CAP. However, despite having observed a positive trend in the use of the PSI and its interpretation, a low-intensity intervention such as the one developed in our work is not sufficient to generalize its use, so other strategies should be considered.

Author contributions: EEZ, MRG & FJRR conceived and design of the study. EEZ & MRG collected data. AGN, BSB, JGB & FJRR contributed in analysis of data. EEZ, MRG & FJRR contributed in analysis and interpretation of data. All the authors have drafting, revising and approved the final version of the manuscript and agreed to be accountable for its contents.

Conflict of interest: The authors declare absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical issues: Including plagiarism, informed consent, data fabrication or falsification and double publication or submission have completely been observed by authors.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Author declarations: The authors confirm that all relevant ethical guidelines have been followed and any necessary IRB and/or ethics committee approvals have been obtained.

References

1. Julián-Jiménez A, Palomo de los Reyes MJ, Parejo Miguez R, Laín-Terés N, Cuenca-Boy R, Lozano-Ancín A (2013) Improved management of community-acquired pneumonia in the emergency department. *Archivos de Bronconeumología*. 49 (6): 230240. doi: 10.1016/j.arbres.2012.12.0082.
2. Esteban-Zubero E, García-Muro C, Alatorre-Jiménez MA, Marín-Medina A, López-García CA, Youssef A, Villeda-Gonzalez (2019) Management of community acquired pneumonia in the emergency room. *Iberoamerican Journal of Medicine*. 1 (1): 3-15. doi: 10.5281/zenodo.3402283.

3. Rider AC, Frazee BW (2018) Community-acquired pneumonia. *Emergency Medicine Clinics of North America*. 36 (4): 665683. doi: 10.1016/j.emc.2018.07.001.
4. Menéndez R, Torres A, Zalacaín R, Aspa J, Martín-Villasclaras JJ, Borderías L, Benítez-Moya JM, et al. (2005) Guidelines for the treatment of community-acquired pneumonia: predictors of adherence and outcome. *American Journal of Respiratory and Critical Care Medicine*. 172 (6): 757762. doi: 10.1164/rccm.200411-1444OC.
5. Labarere J, Stone RA, Obrosky DS, Yealy DM, Meehan TP, Fine JM, Graff LG, Fine MJ (2007) Comparison of outcomes for low-risk outpatients and inpatients with pneumonia: A propensity-adjusted analysis. *Chest*. 131 (2): 480-488. doi: 10.1378/chest.06-1393.
6. Julián-Jiménez A, González-Castillo J, Candel González FJ (2013) When, where and how should a patient with community acquired pneumonia be admitted?. *Revista Clinica Espanola (Barc)*. 213 (2): 99107. doi: 10.1016/j.rce.2012.02.006.
7. Mandell LA, Wunderink RG, Anzueto A, Bartlett JG, Campbell GD, Dean NC, Dowell SF, File Jr TM, Musher DM, Niederman MS, Torres A, Whitney CG (2007) Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clinical Infectious Diseases*. 44 (2): S27-S72. doi: 10.1086/511159.
8. Rodríguez de Castro F (2005) The influence of specialty care on the management of hospitalized pneumonia]. *Archivos de Bronconeumologia*. 41 (6): 297-299. doi: 10.1016/s1579-2129(06)60228-0.
9. González-Moraleja J, Sesma P, González C, López ME, García JF, Alvarez-Sala JL, Cuál es el coste de las neumonías que ingresamos inadecuadamente? (1999) What is the cost of inappropriate admission of pneumonia patients?. *Archivos de Bronconeumologia*. 35 (7): 312-316. doi: 10.1016/s0300-2896(15)30067-3.
10. Ewig S, Torres A (2011) Community-acquired pneumonia as an emergency: time for an aggressive intervention to lower mortality. *European Respiratory Journal*. 38 (2): 253-260. doi: 10.1183/09031936.00199810.
11. Dean NC, Bateman KA, Donnelly SM, Silver MP, Snow GL, Hale D (2006) Improved clinical outcomes with utilization of a community-acquired pneumonia guideline. *Chest*. 130 (3): 794-799. doi: 10.1378/chest.130.3.794.
12. Capelastegui A, España PP, Quintana JM, Gorordo I, Sañudo C, Bilbao A (2006) Evaluation of clinical practice in patients admitted with community-acquired pneumonia over a 4-year period. *Archivos de Bronconeumologia*. 42 (6): 283-289. doi: 10.1016/s1579-2129(06)60144-4.
13. Blasi F, Iori I, Bulfoni A, Corrao S, Costantino S, Legnani D (2008) Can CAP guideline adherence improve patient outcome in internal medicine departments?. *European Respiratory Journal*. 32 (4): 902-910. doi: 10.1183/09031936.00092607.
14. Hu WP, Zhang FY, Zhang J, Hang JQ, Zeng YY, Du CL, Jie Z-J, Jin XY, Zheng C-X, Luo X-M, Huang Y, Cheng Q-J, Qu J-M (2020) Initial diagnosis and management of adult community-acquired pneumonia: a 5-day prospective study in Shanghai. *Journal Thoracic Disease*. 12 (4): 1417-1426. doi: 10.21037/jtd.2020.03.02.
15. Delgado M, Alvarez MM, Carrascosa I, Rodríguez-Velasco M, Barrios JL, Canut A (2013) The routine use of the pneumonia severity index in the emergency department: effect on process-of-care indicators and results in community acquired pneumonia. *Enfermedades Infecciosas y Microbiología Clínica*. 31 (5): 289-297. doi: 10.1016/j.eimc.2012.04.012.
16. Rodriguez JJ, Pastor AJ, Ferré-Llosa C, Bardés-Robles I (2014) Pneumonia severity index in the emergency department and the use of a short-stay unit to improve outcome indicators in community-acquired pneumonia. *Enfermedades Infecciosas y Microbiología Clínica*. 32 (2): 132. doi: 10.1016/j.eimc.2013.07.019.
17. Cabana MD, Rand CS, Powe NR, Wu AW, Wilson MH, Abboud PA, Rubin HR (1999) Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA*. 282 (15): 1458-1465. doi: 10.1001/jama.282.15.1458.
18. Aujesky D, McCausland JB, Whittle J, Obrosky DS, Yealy DM, Fine MJ (2009) Reasons why emergency department providers do not rely on the pneumonia severity index to determine the initial site of treatment for patients with pneumonia. *Clinical Infectious Diseases*. 49 (10): e100-e108. doi: 10.1086/644741.
19. Julián-Jiménez A, Adán Valero I, Beteta López A, Cano Martín LM, Fernández Rodríguez O, Rubio Díaz R (2018) Recommendations for the care of patients with community-acquired pneumonia in the Emergency Department. *Revista Espanola de Quimioterapia*. 31 (2): 186-202.
20. Fine MJ, Auble TE, Yealy DM, Hanusa BH, Weissfeld LA, Singer DE, Coley CM, Marrie TJ, Kapoor WN (1997) A prediction rule to identify low-risk patients with community-acquired pneumonia. *The New England Journal Medicine*. 336 (4): 243-250. doi: 10.1056/NEJM199701233360402.
21. Battleman DS, Callahan M, Thaler HT (2002) Rapid antibiotic delivery and appropriate antibiotic selection reduce length of hospital stay of patients with community-acquired pneumonia: link between quality of care and resource utilization. *Archives of Internal Medicine*. 162 (6): 682-688. doi: 10.1001/archinte.162.6.682.
22. Serisier DJ, Bowler SD (2007) Effect of a simple educational intervention on the hospital management of community-acquired pneumonia. *Respirology*. 12 (3): 389-393. doi: 10.1111/j.1440-1843.2007.01058.x.

23. Halm EA, Horowitz C, Silver A, Fein A, Dlugacz YD, Hirsch B, Chassin MR (2004) Limited impact of a multicenter intervention to improve the quality and efficiency of pneumonia care. *Chest*. 126 (1): 100-107. doi: 10.1378/chest.126.1.100.
24. Yealy DM, Auble TE, Stone RA, Lave JR, Meehan TP, Graff LG, Fine JM, Obrosky DS, Mor MK, Whittle J, Fine MJ (2005) Effect of increasing the intensity of implementing pneumonia guidelines: a randomized, controlled trial. *Annals of Internal Medicine*. 143 (12): 881-894. doi: 10.7326/0003-4819-143-12-200512200-00006.
25. Blasi F, Iori I, Bulfoni A, Corrao S, Costantino S, Legnani D (2008) Can CAP guideline adherence improve patient outcome in internal medicine departments? *European Respiratory Journal*. 32 (4): 902-910. doi: 10.1183/09031936.00092607.
26. Renaud B, Coma E, Labarere J, Hayon J, Roy PM, Boureaux H, Moritz F, Cibien JF, Guerin T, Carre E, Lafontaine A, Bertrand MP, Santin A, Brun-Bussin C, Fine MJ, Roupie E (2007) Routine use of the Pneumonia Severity Index for guiding the site-of-treatment decision of patients with pneumonia in the emergency department: a multicenter, prospective, observational, controlled cohort study. *Clinical Infectious Diseases*. 44 (1): 41-49. doi: 10.1086/509331.
27. Hinojosa Mena-Bernal J, Hinojosa Mena-Bernal C, González Sarmiento E, Almaráz Gómez A, Martín Santos S, Zapatero Gaviria A (2011) Adequacy of the admissions and care provided to the patients with community-acquired pneumonia. *Revista Clinica Espanola*. 211 (4): 179-186. doi: 10.1016/j.rce.2009.06.019.
28. Labarere J, Stone RA, Scott Obrosky D, Yealy DM, Meehan TP, Auble TE, Fine JM, Graff LG, Fine MJ (2006) Factors associated with the hospitalization of low-risk patients with community-acquired pneumonia in a cluster-randomized trial. *Journal of General Internal Medicine*. 21 (7): 745-752. doi: 10.1111/j.1525-1497.2006.00510.x.
29. Calbo E, Ochoa de Echagüen A, Rodríguez-Carballeira M, Ferrer C, Garau J (2004) Hospital admission, duration of stay and mortality in community-acquired pneumonia in an acute care hospital. Correlation between a pneumonia prognosis index and conventional clinical criteria for assessing severity. *Enfermedades Infecciosas y Microbiología Clinica*. 22 (2): 64-69. doi: 10.1016/s0213-005x(04)73036-0.
30. Riccioni G, Di Pietro V, Staniscia T, De Feudis L, Traisci G, Capani F, Ferrara G, Di Ilio E, Di Tano G, D'Orazio N (2005) Community acquired pneumonia in internal medicine: a one-year retrospective study based on pneumonia severity index. *International Journal of Immunopathology and Pharmacology*. 18 (3): 575-586. doi: 10.1177/039463200501800318.